

# *Mathematical Journal of Okayama University*

---

*Volume 14, Issue 1*

1969

*Article 3*

NOVEMBER 1969

---

## A note on separable extensions of commutative rings

Motoshi Hongan<sup>\*</sup>

Takasi Nagahara<sup>†</sup>

<sup>\*</sup>Okayama University

<sup>†</sup>Okayama University

Copyright ©1969 by the authors. *Mathematical Journal of Okayama University* is produced by  
The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

## A NOTE ON SEPARABLE EXTENSIONS OF COMMUTATIVE RINGS

MOTOSHI HONGAN and TAKASI NAGAHARA

Commutative separable algebras have been studied in [1] and [2], [3] where the main ideas are based on the theory of fields. Moreover, in [4], G. J. Janusz presented a number of explicit results for commutative separable algebras. In this paper, we shall make a remark on commutative separable algebras (Theorem), where this paper depends heavily on [4].

Throughout the present paper,  $A$  will be a commutative ring with the identity element 1, and  $B$  a subring of  $A$  containing the identity element 1 of  $A$ . As in [4], if  $A$  is projective, and separable over  $B$  then  $A$  will be called a strongly separable  $B$ -algebra. By [5, Villamayor's Theorem], a strongly separable  $B$ -algebra is a finitely generated  $B$ -module. Moreover, by [4, Th. 1.1], a strongly separable  $B$ -algebra with no proper idempotents (no idempotents except 0 and 1) can be imbedded in a Galois extension of  $B$  with no proper idempotents. If  $N$  is a Galois extension of  $B$  then the Galois group will be denoted by  $G(N/B)$ . Our purpose of this paper is to prove the following theorem.

**Theorem.** *Let  $A \supseteq B$ , and  $A$  a strongly separable  $B$ -algebra without proper idempotents. Then,  $B[a]$  is separable over  $B$  for every  $a \in A$  if and only if  $A$  is a field.*

Firstly, we shall prove the following

**Lemma.** *Let  $B[a]$  be a commutative ring without proper idempotents, and  $a \notin B$ . If  $B[a]$  is a strongly separable  $B$ -algebra then  $a$  is not nilpotent.*

*Proof.* Let  $N$  be a ring extension of  $B[a]$  without proper idempotents which is Galois over  $B$ . Then, there exists an element  $\sigma$  of  $G(N/B)$  such that  $a \neq a^\sigma = b$ . By [4, Lemma 2.7 and Lemma 2.1],  $a - b$  is an invertible element of  $N$ . If  $a^n = 0$  for some natural number  $n$  then  $b^n = 0$ , and so, we have a contradiction

$$0 \neq (a-b)^{2n} = \sum_{i=0}^{2n} (-1)^i \binom{2n}{i} a^{2n-i} b^i = 0$$

Hence  $a$  is not nilpotent.

Now, we shall prove our theorem.

*The proof of Theorem.* Let  $N$  be a ring extension of  $A$  without proper idempotents which is Galois over  $B$ . If  $A$  is a field then, for every nonzero element  $a$  of  $B$ , we have  $(a^{-1})^\sigma = (a^\sigma)^{-1} = a^{-1}$  for all  $\sigma \in G(N/B)$ , which implies  $a^{-1} \in B$ ; hence  $B$  is a field, and  $A$  is separable over  $B$  in the sense of classical separability, and so, every element of  $A$  is separable over  $B$ . Conversely, we assume that  $B[a]$  is separable over  $B$  for every  $a \in A$ . At first, we shall prove that  $B$  is a field. Let  $a$  be an element of  $A$  which is not contained in  $B$ . Then, there exists an element  $\sigma$  of  $G(N/B)$  such that  $a \neq a^\sigma$ . By [4, Lemma 2.7 and Lemma 2.1],  $a - a^\sigma$  is an invertible element of  $N$ . Let  $b$  be a nonzero element of  $B$ . Then  $b(a - a^\sigma) \neq 0$ . Hence  $ba \neq (ba)^\sigma$ , and so,  $ba - (ba)^\sigma = b(a - a^\sigma)$  is an invertible element of  $N$ . Therefore  $b$  is an invertible element of  $N$ . Since  $(b^{-1})^\tau = (b^\tau)^{-1} = b^{-1}$  for all  $\tau \in G(N/B)$ , we have  $b^{-1} \in B$ . Thus  $B$  is a field. Since  $A$  is a finitely generated  $B$ -module,  $A$  is an Artinian ring. By lemma, the radical of  $A$  is zero. Hence  $A$  is a semisimple ring. Noting that  $A$  has no proper idempotents,  $A$  is a field.

The following corollaries are direct consequences of our theorem.

**Corollary 1.** *Let  $A \not\cong B$ , and  $A$  a strongly separable  $B$ -algebra without proper idempotents. If  $A$  is not a field then there exists an intermediate ring  $B[a]$  of  $A/B$  such that  $B[a]$  is not separable over  $B$ .*

**Corollary 2.** *Let  $A$  be a strongly separable  $B$ -algebra without proper idempotents. If  $B[a]$  is separable over  $B$  for every  $a \in A$  then, for every intermediate ring  $D$  of  $A/B$ ,  $D = B[d]$  for some element  $d$  of  $A$ .*

**Corollary 3.** *Let  $A \not\cong B$  and suppose  $A$  has no proper idempotents. Then,  $B[a]$  is strongly separable over  $B$  for every  $a \in A$  if and only if  $A$  is a field which is algebraic and separable over  $B$ .*

REFERENCES

- [1] M. AUSLANDER and D. BUCHSBAUM: On ramification theory in Noetherian rings, Amer. J. Math. 81 (1959), 749—765.
- [2] M. AUSLANDER and O. GOLDMAN: The Brauer group of a commutative ring, Trans. Amer. Math. Soc. 97 (1960), 367—409.
- [3] S. CHASE, D. K. HARRISON and A. ROSENBERG: Galois theory and cohomology of commutative rings, Mem. Amer. Math. Soc. No. 52 (1965).
- [4] G. J. JANUSZ: Separable algebras over commutative rings, Trans. Amer. Math. Soc. 122 (1966), 461—479.
- [5] T. NAGAHARA: A note on Galois theory of commutative rings, Proc. Amer. Math. Soc. 18 (1967), 334—340.

DEPARTMENT OF MATHEMATICS,  
OKAYAMA UNIVERSITY

*(Received March 31, 1969)*